

Climate Change & Commercial Fisheries in Point Judith, RI

INTRODUCTION




Climate change is altering the physical and chemical characteristics of our ocean and affecting marine ecosystems and fisheries. As environmental conditions continue to change, fishing communities may be affected by changes in the distribution and availability of species. This report summarizes the current status of fisheries in Point Judith and shares information on changes in harvested species that may occur in the future. Used alongside the [Climate Adaptation Resource Hub for Fishing Communities](#), this report provides information for understanding potential impacts on a fishing community, which can be used to consider ways to adapt to a changing climate.

WHAT IS LANDED HERE?

A diverse mix of commercially harvested species are landed in Point Judith, representing an annual average of 46.5 million pounds valued at \$55.1 million from 2012-2021.* Longfin squid was the dominant species landed in the port, contributing an annual average 11.6 million pounds, valued at \$14.7 million on average per year. Sea scallops also made substantial contributions, with landings averaging 1.3 million pounds per year, valued at an average of \$13.3 million per year. The total volume of landings in Point Judith experienced a decline from 1998 until 2011. Landings rose to 54 million pounds in 2015 and have declined in recent years. The total value of landings has fluctuated over the years but is generally increasing.

Species	Annual Average Value	Annual Average Volume
Longfin squid	\$14,709,941	11,597,016 lbs
Sea scallop	\$13,340,777	1,276,174 lbs
American lobster	\$5,912,551	1,124,353 lbs
Summer flounder	\$5,018,527	1,488,959 lbs
Scup	\$2,804,663	4,949,437 lbs
Silver hake	\$2,071,215	3,411,818 lbs
Jonah crab	\$1,161,944	1,476,550 lbs
Other	\$10,098,200	21,154,528 lbs


Above are the annual average value and volume for the top species landed at this port in each year from 2012-2021.



*Landings data were provided by NOAA Fisheries' Greater Atlantic Regional Fisheries Office. Due to confidentiality restrictions, some data may not be fully representative of the historical landings at a given location.


LANDINGS OVER TIME

Lobster landings in Point Judith have been generally declining since the early 2000s. Herring landings have varied and fallen to low levels in recent years. Longfin squid landings have fluctuated around 10 million pounds, with several intermittent years of high landings. Scup landings increased throughout the 2000s, plateaued in the mid-2010s and have declined since. Summer flounder has fluctuated around 1.5 million pounds, with periods of high landings in 2002-2006 and 2010-2016 and a recent uptick since 2019. Silver hake landings declined through much of the 2000s and 2010s but have recently increased.




OUR CHANGING CLIMATE AND WARMING WATERS

Greenhouse gas emissions around the world are a primary contributor to the warming the planet has been experiencing over the past century. This warming affects the health and distribution of species that support fisheries in coastal communities. Scientists around the world use a common set of scenarios to project climate impacts into the future. These scenarios represent multiple global social and economic development patterns paired with different levels of greenhouse gases in Earth's atmosphere. The scenario representing the largest build-up of greenhouse gases, labeled SSP5-8.5, indicates global average temperatures will warm by approximately 4°C (7°F) above pre-industrial levels by the end of this century. We use this scenario to understand how species may respond to changes in ocean temperatures in the Northeast U.S. relative to those experienced during 2010-2019. These species projections allow us to explore different potential futures of fisheries and support decisions now that can buffer the severity of future climate change impacts on fishing communities.



FUTURE CHANGES IN AVAILABILITY

As the abundance and distribution of certain species changes with warming waters, communities may need to respond to ensure the continuity of the fishing industry. By combining historical species observations with future climate information, we can estimate how the availability of certain species may change, and what new opportunities may emerge. Availability is given here as the total estimated weight of a particular species of fish in a given area, as modeled from bottom trawl survey data. Warming ocean temperatures may affect the availability of some commercial species in the waters near Point Judith. Black sea bass, rock crab, and summer flounder may increase with increasing temperatures. Longfin squid and herring availability may vary with different levels of warming. Lobster and silver hake may increase at lower levels of warming but decline slightly at higher temperatures; conversely, scup may decline with low levels of warming but increase as warming levels rise.



EMERGING OPPORTUNITIES AND ADAPTATION OPTIONS

Harvesting emerging species and diversifying catch are some ways individual harvesters can adapt to changing fisheries. In the table below, we outline other potential adaptation options spanning the different scales of the fishery system. As the climate continues to change, new impacts will take shape, requiring re-evaluation and revision of goals in order to respond to climate change. For more information on adaptation options in fishing communities, please visit the [Climate Adaptation Resource Hub for Fishing Communities](#).

Individual Harvester Actions	Industry Actions
<ul style="list-style-type: none"> • Shifting fishing locations • Shifting harvested species • Diversifying livelihood (alternative fisheries, aquaculture, non-fishing jobs) 	<ul style="list-style-type: none"> • Improving product handling • Developing supply chain capacity • Diversifying markets and building consumer demand
Management Measures	Community Initiatives
<ul style="list-style-type: none"> • Reassessing quota allocations • Altering permit access and availability • Developing adaptive reference points • Applying dynamic and ecosystem-based management 	<ul style="list-style-type: none"> • Maintaining and securing shoreside infrastructure • Improving transportation networks • Developing local seafood initiatives • Conducting vulnerability and resilience assessments • Using early warning monitoring • Community adaptation and resilience planning

Projected Changes in Species Availability in Point Judith

Values represent percent change in modeled species availability at potential levels of warming relative to 2010–2019 baseline conditions. Species in gray had low availability (<5 lbs/sq. mile) during the baseline period.

Species	Increase in Sea Surface Temperature			
	0.9°F	1.8°F	3.6°F	5.4°F
Acadian redfish	-2.3%	-4.7%	-29.9%	-45.5%
American lobster	8.4%	22.8%	14.8%	-2.6%
American plaice	-4.5%	-5.5%	-13.6%	-18.1%
Atlantic cod	-1.0%	37.5%	54.1%	34.3%
Atlantic halibut	5.4%	2.7%	1.4%	-6.7%
Atlantic herring	16.0%	8.5%	-21.3%	7.5%
Atlantic mackerel	24.3%	44.0%	79.9%	68.9%
Black sea bass	29.8%	89.0%	108.6%	345.3%
Butterfish	2.9%	-11.3%	18.0%	2.7%
Deep sea red crab	-25.4%	-22.6%	-38.3%	-28.7%
Haddock	1.3%	25.6%	24.7%	-28.2%
Hagfish	42.8%	97.6%	96.1%	39.6%
Jonah crab	15.4%	-6.9%	-1.4%	-31.5%
Little skate	-41.9%	-41.8%	-37.1%	-32.9%
Longfin squid	45.1%	60.8%	-13.2%	149.1%
Monkfish	-12.0%	-18.8%	-33.5%	-41.0%
Ocean quahog clam	-37.0%	-7.3%	32.2%	-47.1%
Pollock	-14.6%	2.3%	-5.4%	-19.7%
Red hake	34.2%	25.7%	7.8%	-1.9%
Rock crab	70.1%	71.4%	111.9%	184.8%
Sand lance	-2.6%	-9.0%	-28.1%	-28.5%
Scup	-45.6%	-27.4%	14.4%	108.0%
Sea scallop	21.0%	15.0%	-19.8%	-22.6%
Shortfin squid	-44.9%	-22.4%	-32.1%	-16.8%
Silver hake	5.1%	7.4%	-0.8%	-15.3%
Smooth skate	-20.8%	-21.3%	-15.7%	-15.1%
Spiny dogfish	-72.9%	-71.0%	-70.4%	-72.1%
Summer flounder	1.1%	25.1%	56.9%	113.4%
Thorny skate	-21.2%	-15.6%	-22.2%	-35.5%
White hake	-17.3%	-8.7%	3.8%	5.2%
Windowpane	4.7%	29.2%	21.4%	39.5%
Winter flounder	8.7%	11.0%	14.3%	3.7%
Winter skate	-47.9%	-51.3%	-42.1%	-35.3%
Witch flounder	-3.8%	-26.0%	-31.7%	-29.6%
Yellowtail flounder	-32.1%	-36.6%	-47.6%	-42.1%

MAKING SENSE OF CLIMATE PROJECTIONS AND SPECIES DISTRIBUTION MODELS

The species results shown here were developed using a spatio-temporal species distribution model, which can estimate the current and future distribution of marine species through time and space. The model uses projected regional sea surface and bottom temperature data from the globally coordinated Coupled Model Intercomparison Project (CMIP6) and species data from bottom trawl surveys conducted by the Northeast Fisheries Science Center and the Department of Fisheries and Oceans. Estimated species biomass densities are then averaged over an area fished by vessels from the port of interest. This enables us to interpret local changes in availability of a species at a specific time or temperature.

LEARN MORE

For more information regarding climate change, species distribution change, fisheries adaptation options, and adaptation barriers and enablers, please visit:

gmri.org/adaptationhub

ASK QUESTIONS

For specific questions regarding your community, contact Kathy Mills at:

kmills@gmri.org



**Gulf of Maine
Research Institute**

Science. Education. Community.